

CASE REPORT

By

Dr. Selig B. Kousnetz, O.D., M.C.S.O.

Pseudo Myopia

Mrs. E.G., age 19, Type P.

She complains that distance vision is blurred; she has worn glasses for 8 years, and had them changed about 4 times.

Present Rx by medical eye specialist, about one year old, O.U. -1.75 with visual acuity, of Right, left and O.U. 20/20.

Uncorrected vision, Right 20/30    Left 20/30    OU 20/25-2

Health, excellent.

Versions – rotations – movement of eyes, good.

Absolute convergence, 2 1/2".

Pupils 5 MM. Pupillary reflexes, good and bi-consensual. Iris – lids- conjunctiva – lachrymal apparatus, normal. Cornea and lenses, clear.

- (1) Ophthalmoscope neg. PD 6/63
- (3) Habitual phoria distance ½ EXO.
- (4) Static retinoscope OU - .50.
- (5) Dynamic Retinoscope at 20" OU +1.00.
- (6) Dynamic Retinoscope at 40" OU Plano.
- (7) Subjective OU -.50. Visual acuity, OU 20/20.
- (8) Induced phoria ½ EXO
- (9) True adduction, 7.
- (10) Convergence 24/16.
- (11) Abduction 9/7
- (12) Vertical phoria-negative.
- (13B) Induced phoria at near 2 eso.
- (14A) Cross cylinder dissociating OU +.25.
- (15A) Induced phoria 3 es0.
- (16A) Total adductive stimulation 19.
- (17A) Total adductive inhibition 2 exo.
- (17B) Negative fusional reserve 1/-4.
- (18) Vertical phoria a near, negative.



