

MINIMUM ROUTINE FOR CASE REPORTS OF PATIENTS

1. Old Rx. Visual acuity
2. Naked Vision
3. Amplitude of accommodation
4. Phoria, old or naked, distance
5. Infinity, P. D.
6. Duction, Infinity, new Rx
7. Shock test distance Rx
8. Phoria near, new Rx, 13"
9. Duction, near point, new Rx 13"
10. Amplitude of accommodation with new Rx
11. Shock test near point with new Rx, 13"

1. Old Rx Visual acuity
Obvious

2. Naked vision
Obvious

3. Amplitude of accommodation
Amplitude of accommodation with old Rx or no Rx if never worn, by the approved method and record in dioptric equivalent of blur out point using printer & 8-point type. Binocular.

In presbyopes lend +2.00 S

Same technique for 10 amplitude of accommodation with new Rx.

4. Phoria, old Rx or naked, distance
Maddox double prism
Maddox rod
Maddox multiple rod if this rod is an instrument reverse 180 degrees and make a second test.

Test object must be below level of eyes.

MINIMUM ROUTINE FOR CASE REPORTS OF PATIENTS

5. Infinity P.D.

Arm's length from patient who looks at your pupil while other eye is closed, then alternate to next eye and patient looks at that pupil.

6. Duction, Infinity, new Rx

Base-in to break and recovery, and then base-out to break and recovery with spot as test object not more than 1 cm. in diameter at 20' and below the level of the eyes.

7. Shock fusion test, distance, new Rx

One cm. spot at infinity cover both eyes, remove card, replace, turn base-in prism with card before eyes. Remove card, check for diplopia. Increase prism power while eyes are covered with card. Continue until cannot fuse. The last power than can fuse is measure to record.

8. Phoria, near, new Rx 13"

Phoria at near. Disassociation method with black spot not over 3 mm. in diameter and not using more than 5Δ up and 5Δ down. Or use the Maddox rod and spot of light not more than ½ mm. in diameter.

9. Duction, near point, new Rx 13"

Base-in to blur-out, break, recovery.

Base-out to break and recovery. Use vertical row of 8 point periods as test object, or letters not larger than those used for the amplitude test. 13"

10. Amplitude of accommodation with new Rx.

Same technique as for No. 3 with new Rx.

11. Shock test, near point, new Rx, 13"

Same technique at near, 13", as No. 7, using same test object as used in duction test at near, vertical row of 8 point periods, or letters not larger than these used for amplitude of accommodation.