



THE PUPIL

Observation and Grading

Alpha Omega Pupil

John J. Pulaski, O.D., FCSO

College of Syntonic Optometry

101 Course

Cambridge, Canada Jan 2025

The Pupil And Syntonics

How do you know if a person
needs Syntonic Treatment?

The Pupil And Syntonics

Three keys elements in Syntonic Clinical
evaluation and treatment application.

1. The Pupil - AO
2. The Field - Kinetic
3. The Patient History

The Pupil

One of the most sensitive measures of ANS activity.

- Neurologic marker of ANS/Midbrain/Central brain function.
- “Eyes are the Window to the Soul”
The Pupils are the Windows.
- Portal of Energy Reception and Projection. Portal through which we interact with our world.
- Non-verbal Communication and strong emotional indicator.
- Reception of “nutrition” - LIGHT



The Pupil

Neurological Pathways

Sphincter Pupillae

LIGHT

Pretectal Nucleus

I
Optic Nerve (II)
(Afferent Arm)

Retina

Oculomotor Nerve (III)
(Efferent Arm)

EXCITATION

Ciliary Ganglion

Pupil

E.W.

Parasympathetic Pathway

INHIBITION

Sympathetic Pathway


Opioids Block

I.N.

Dilator Iridis


From Superior Cervical Ganglion

STRESS
(Hypercarbia, Hypoxia,
and Nociception)



The Pupillary Light Reflex as a Biomarker of Concussion

Frederick Robert Carrick^{1,2,3,4,5}, [Sergio F Azzolino](#)⁵, [Melissa Hunfalvy](#)⁵, [Guido Pagnacco](#)^{5,6}, [Elena Oggero](#)^{5,6}, [Ryan C N D'Arcy](#)^{7,8,9}, [Mahera Abdulrahman](#)¹⁰, [Kiminobu Sugaya](#)^{1,2}



The Pupil

Neurological Pathways



Parasympathetic - Constriction

- The Pupillary Light Reflex (PLR)
- Influence on Iris Sphincter
- Light-Inhibited Sympathetic Path
- Trigeminal Nerve – sensory stimulation to eye/iris

Sympathetic – Dilation

- Direct stimulation of Iris Dilator through 3 neuron arc
- Inhibition of EW nucleus

Pulaski AO Pupil




The Pupil

Neurological Pathways

Pathway of Dilation

- Direct Stimulatory - Iris dilator
 - Hypothalamus
 - Nucleus Coeruleus
 - Superior Colliculus
 - Frontal Eye Fields
- Inhibitory – EW nucleus
 - Nucleus Coeruleus
 - Superior Colliculus
 - Supernuclear Inhibition from the Reticular Activating Formation in brainstem



The Pupil In Syntonics

In Syntonics we are interested in the pupil as an indication of the well being of our patient particularly in regards to the state of the ANS. The two major reactions observed are:

- Alpha Omega Pupil
- Pupillary Light Reflex



The Pupil

Reactions

Alpha Omega Pupil



- An Alpha Omega Pupil is the abnormal re-dilation of the pupil during direct, constant light stimulation.
- Unique to the practice of Syntonics
- First suggested as a term by Dr. Paul Johnson in 1934.
- The abnormality is brought to normalcy with phototherapy treatment
- There is an inverse relation between the size of the functional visual field and the degree of the AO Pupil.



4.5 to 2.75 4+PLR



6.5 / 4.22 2-3+ PLR, 5 then 5.25, 2nd time 6 /4.6 to 5.5,

THE NORMAL PUPIL

THE PUPIL





Anatomy, Physiology, and Clinical Applications

IRENE E. LOEWENFELD, Ph.D.

Based on the cooperative work with
Otto Lowenstein, M.D.



Pupillary Reactions




Latency	Time from light flash to pupil constriction
MaxPD	The maximum diameter of pupil before light flash
MinPD	The minimum diameter of pupil after light flash
MCV	The maximum constriction velocity of pupil constriction after light flash
75% PRT	The time for redilation of pupil to 75% of its previous maximum diameter after light flash

The Pupil

The “Normal” Pupil

- Steady state size
- Dynamic, transient response
- Neurons in E-W Nucleus generate steady firing rate even when no external inputs are involved. Balanced by cortical inhibition.
- Slight latent period before contracts
- Maximum contraction is reached in about 2 sec.
- When stimulus removed pupil re-dilates much slower than it contracted – Recovery response
- Under prolonged stimulation there are normal oscillations


Pupillary Reactions



Pupillary Reactions

Bright Lamp

Carrick et al Life (Basel). 2021 Oct; 11(10): pii: 10.3390/life11101104



Latency	Time from light flash to pupil constriction
MaxPD	The maximum diameter of pupil before light flash
MinPD	The minimum diameter of pupil after light flash
MCV	The maximum constriction velocity of pupil constriction after light flash
75% PRT	The time for redilation of pupil to 75% of its previous maximum diameter after light flash

The Pupil

Pupillary Light Reflex

MCA, MCV best predictors for
Parasympathetic Dysfunction

Alzheimer's

Parkinson's


ABI/TBI

Autism

Drugs/Alcohol

Infection

ADD – increased MPD and MCV



The Pupil



Reactions

Pupillary Light Reflex

- 10-20% of RGC axons relate to PLR
- 1% of ipRGC's
- Speed/Strength of contraction
- Coma, TBI, Stroke
- Excited Patient
- Relation to ANS

Pulaski AO Pupil

Pupillary Light reflex




The Pupil


Reactions

Pupillary Light Reflex

- ipRGC's very important in recovery phase (PIPR)
- Pupil both inhibited and activated by different wavelength sensitive cones
- Important in sustained constriction



Synaptic Input/Output ipRGC



The Pupil

The "Normal" Pupil

Pupillogram – 24 y.o. healthy male

•From Loewenfeld, "The Pupil"




Figure 10-1. Spontaneous pupillary movements in a healthy 24-year-old man. The pupillograms of the right eye are shown. A and C were original records. In B, the time axis of the graph was compressed in the following manner: the pupillary diameter at the beginning of each successive second was taken from the original record, and was plotted as a single measurement (small rectangles); sixty such measurements thus show the pupillary movements

that occurred within each minute. A: Normal light reflex; B: Diameter during the fifth to ninth minute in darkness. The pupil was large and quiet. The subject was not tired, and was able to continue the test for more than 2 hours without remarkable changes in pupillary behavior. C: Slow (a) and fast (b) pupillary oscillations of small extent (see text). (From O. Lowenstein, R. Feinberg, and I.E. Loewenfeld, *Invest. Ophthalmol.*, 2 [1963]:138)

The Pupil

The “Normal” Pupil

- Pupillary Unrest
- Pupillary Fatigue
- Pupillary Escape

The Pupil

The “Normal” Pupil Pupillary Unrest

- Defined by Lowenfeld as “normal pupillary oscillations brought on by steady light and absent in darkness.”
- They are continuous, constant rippling and pumping, rapid and irregular movements of the pupil in a lighted environment that cease when the light is turned off.
- Can vary from one individual to another but the pattern is the same for that individual over time. Postulated that it is unique due to our genetic make-up.
- Their cause remains unknown although it is Lowenfeld’s opinion that they originate in the neurons of the midbrain.

The “Normal” Pupil

Pupillary Unrest

Pupillogram of normal 24 y.o. female under sustained light (Loewenfeld, “The Pupil”)




Figure 3-67. Pupillary oscillations during unilateral adaptation to a bright, steady light. Pupillogram of a normal 24-year-old woman. The solid line represents the right, the broken line the left pupil. **First line:** The pupils were large and quiet in darkness. When the right eye was exposed to a steady, bright light, both pupils contracted, then redilated somewhat and began to oscillate. **Second line:** Pupillary oscillations after the right eye had been

light was turned off (arrow), the pupils enlarged and the oscillations disappeared. The movements of the right and left pupils remained equal throughout the experiment, even though the right eye alone was stimulated, while the left eye remained in darkness. (From O. Lowenstein and I.E. Loewenfeld, *Amer. J. Ophthalmol.*, 48, II [1959]:536; published with permission of *The American Journal of Ophthalmology*, ©The Ophthalmic Publishing Company)



OD 6.5 /4.5 to 4.64 to 5, OS 6.5 /4.5 to 4.6

The Pupil

The “Normal” Pupil

Fatigue Waves

- They originate from fluctuation in central nervous system integration.
- They begin to occur with a decrease in sympathetic discharges and inhibitory influences on the EW nucleus causing increased constriction of the pupil.
- Causes include boredom, sleepiness, illness and ageing.
- They decrease during periods of arousal, when the person is mentally active.
- The reflex becomes well integrated after a good night's rest.
- **Adrenal Fatigue**

The Pupil

The “Normal” Pupil

Fatigue Waves




Figure 10-2. Pupillary “fatigue waves.” In each line, the diameter of the subject’s right pupil was plotted as the ordinate by the same method described for Figure 10-1. **A:** Record obtained from a normal, alert 24-year-old man. The pupils showed only little activity after many minutes in darkness. The graph shows the movements between the twelfth and sixteenth minute of the experiment. **B to D:** Pupillary movements of a healthy but very tired 38-year-old man. Less than 3 minutes after the beginning of the test the pupils began to become smaller. During the following minutes extensive, irregular waves of pupillary contraction and dilation appeared as the subject repeatedly drifted toward sleep and roused himself spontaneously (lines **B** and **C**). Finally, the periods of arousal became shorter and shorter and less and less complete. Immediately before he fell asleep, the pupils were very small (end of line **D**). (From O. Lowenstein and I.E. Loewenfeld, *Annls N.Y. Acad. Sci.*, 117 [1964]:142)

•From Loewenfeld, “The Pupil”
Pulaski AO Pupil








Figure 2-1. Spontaneous pupillary movements in darkness. **A** and **B**: Pupillary behavior of two healthy young men during $3\frac{1}{2}$ minutes in darkness on four different days, at intervals of 1 week (lines 1 to 4). The records were taken between 5 and 6 P.M. Both subjects were conscientious dental students who had worked hard in the clinic all afternoon. The first subject (**A**) quickly became sleepy as he sat in the dark. The second (**B**) did not. The records show pupillary diameter (in mm) plotted against time (in minutes). They were obtained by contracting the usual time axis of the pupillograms to one-twentieth of its usual length, by measuring the pupillary size at the beginning of each successive second of the original record, and plotting each measurement as a small square on the graph. Subject A's pupils became unsteady already within the first minute of the experiment. In contrast, subject B's pupils never showed more than the slightest variations. The longest test run on B was 90 minutes, but even then his pupils did not waver. **C¹** and **C²**: Light reflexes of the students, elicited by ten consecutive 1-second bright light flashes, with 2-minute rest periods interposed (white light, 5° retinal area, centrally fixated, 7.5 log units above the subjects' scotopic visual threshold). The individual reflexes were superimposed. Note the marked variability in reactions for **A** (**C¹**), and the tight bundle of curves for **B** (**C²**).

The Pupil Reactions Pupillary Escape



- Re-dilation of the pupil after exposure to a moderate light source.
- Stated that the re-dilation is due to retinal adaptation
- Pupillary “Unrest” follows
- The Alpha Omega pupil might be considered as a form of pupillary escape by non-light therapists




The Pupil Reactions Pupillary Escape



- Duke-Elder,¹ in 1959 discussing the clinical picture of optic neuritis, mentions "a peculiar pupillary reaction common to all forms of conduction interference, wherein, although both the direct and consensual reactions are present the contraction is not maintained under bright illumination so that the pupil slowly dilates again while the light is still kept upon the eye."
- "**Pupillary escape**" is an abnormal **pupillary** response to a bright light, in which the **pupil** initially constricts to light and then slowly redilates to its original size. **Pupillary escape** can occur on the side of a diseased optic nerve or retina, most often in patients with a central field defect.

The Pupil Reactions


Pupillary Escape



The Pupil

Reactions

Alpha Omega Pupil



- An Alpha Omega Pupil is the abnormal re-dilation of the pupil during direct, constant light stimulation.
- An Alpha Omega Pupil differs from Pupillary Unrest in that its occurrence happens before 8-9 seconds have elapsed. Its amplitude is greater than 1 mm in many cases.
- It's measurement or recording can be effected by the influence of "Fatigue Waves" in a very tired person and effect the variability of response during testing.
- It can be considered a form of pupillary "escape" but without pathology that is reversible with treatment. There are also differences in frequency and amplitude.
- Conclusion – The pupil reaction is extremely complex

The Pupil

Sympathetic

Influences on Pupillary Reflex Dilation

The Normal Pupillary Reflex Dilation

Any sensory, emotional, or mental stimulus elicits reflex dilation. Any sound, touch or pain, fear, joy or anger or spontaneous thoughts and intentional efforts all dilate the pupils.

The amplitude of reaction depends on the degree of arousal caused by the stimulus and the subject's physical and mental state at the time of stimulation.

Loewenfeld

The Pupil





Influences on Pupillary Reflex Dilation

Increased Dilation

Increased attention
Orienting Reflex
Increased mental effort
Mental arithmetic
Memory formation
Pleasant sounds
Perception of odors
Sexual arousal
Dishonesty, Lying
Enjoy/Dislike what is seen

Increased Constriction

Mind wandering/Distraction
Introspection
Poor task performance
Disgust
Images of the sun
High level scene processing
Memory retrieval



Average change in pupil size in response to visual cues as measured by a pupillometer among 12 non-smokers, and 7 smokers who averaged 10.6 cigarettes per day, after 3 hours without smoking. Chae Y. et al, Subjective and Autonomic Responses to Smoking-Related Visual Cues, The Journal of Physiological Sciences, March 25, 2008 - Graphic compiled using study figures by WhyQuit.com

The Pupil

REACTIONS

The Pupil Reactions

- Alpha Omega
- Hippus
- Afferent Pupillary Defect
- Pupillary Light Response (PLR)
- Near Reflex

The Pupil


Reactions

Alpha Omega Pupil



- Why is it not Hippus?
- Why is it not an APD defect?
- How do other reactions relate?
- How does the field relate?

The Pupil Reactions Alpha Omega Pupil



- An Alpha Omega Pupil is the abnormal re-dilation of the pupil during direct, constant light stimulation.
- It is related to imbalances within the ANS. Hypothalamus and Adrenal activity are important.
- The reaction reveals many aspects of the vitality of the human being.
- Rian Shah, N.D. states that oscillations are due to severe sodium depletion secondary to decreased adrenal function, specifically diminished aldosterone. Also called “rebound” pupil.

The Pupil

Testing Standards in Measurement

Alpha Omega Pupil

Pulaski, 2006

Observation and Recording of AO Pupil

1. Quickness of initial stimulation – PLR
2. Time to release
3. Amplitude of release
4. Reactions after initial release – fluctuations
5. Change in response with repeated stimulation
6. Sensory reactions – tearing, pain, etc

The Pupil

Grading Standardization

Alpha Omega Pupil

Alpha Omega Grading Standard
(Pulaski 2010)

Grade	Release Time	Fluctuations	Amplitude
Normal	≥ 9 seconds	Trace	Trace
1+ AO	4 – 6 sec	Moderate	Mild
2+ AO	2 – 3 sec	Marked	Mild-Moderate
3+ AO	1 – 2 sec	Mild-Moderate	Moderate
4+ AO	< 1 sec	Mild	Large

The Pupil

Grading Summary

OD						OS				
0	1	2	3	4		0	1	2	3	4
0	1	2	3	4	PLR	0	1	2	3	4
					Near Reflex	0	1	2	3	4
Yes	No				Normal Direct	Yes	No			
Yes	No				Normal Consensual	Yes	No			
_____					Pupillary Diameter	_____				
_____					Time of Release	_____				
_____					Amplitude of Release	_____				
0	1	2	3	4	Fluctuations	0	1	2	3	4
0	1	2	3	4	AO Pupil	0	1	2	3	4
_____					Change in repeated stimulation	_____				
_____					Sensory reactions	_____				

Field / Progress Evaluation

DATE ___/___/___ NAME _____ Filter AO/MD MU UO/MU _____

Copay _____ Total Weeks _____ Since LastOV _____

HX Compliance + - _____ Changes in Health/Medication None _____

Field Testing Tested by KL JP Reliability 1 2 3 4 _____

V A sc cc OD 20 _____ Pupils Size SS/Lt PLR $\alpha\Omega$ APD Near NPC _____/_____

OS 20 _____ OD _____/_____ mm 1 2 3 4+ 0 1 2 3 4+ - + + -

OS _____/_____ mm 1 2 3 4+ 0 1 2 3 4+ - + + -

Photophobia None 1 2 3 4+ Other _____

Fields White Green Red Blue BS EOM's Pursuits Saccades Hypo/Hypermtric

OD _____% _____% _____% _____% _____ SAFE 1 2 3 4+ _____% Rt 1 2 3 4+ _____

OS _____% _____% _____% _____% _____ SAFE 1 2 3 4+ _____% Lt 1 2 3 4+ _____

Jerky 1 2 3 4+ Fixation Loss 1 2 3 4+ Sac Int _____+

Dynamic OD _____ Impression ☐ Visual Field Contraction Plan ☐ Continue Current Filters

OS _____ ☐ Improving W C ☐ Same W C ☐ FUp _____ Weeks

☐ Worse OD OS ☐ Asymmetric

☐ Pursuit / Saccadic Dysfunction ☐ Change Tx _____

☐ Pupillary Abnormality _____


☐ Convergence Insufficiency _____

Externals OD OS

Lid/Lash _____

Conjunctiva _____

Signed _____



OD 7 / 4.25 then 5.0 2+ PLR, 1-2+ 2nd
OS 6.0 / 4.5 to 5.0 Rate 2+



6.5 / 4.22 2-3+ PLR, 5 then 5.25, 2nd time 6 /4.6 to 5.5,

The Pupil Reactions

Hippus


Per Lowenfeld it is a very confusing term

- Nystagmus in ancient times
- Lid fluttering
- 18th Century – wave-like pupillary movements
- Nine categories (see addendum – pupillary oscillations)
- Loewenfeld suggests that the name be dropped or defined as “vigorous pupil oscillations of various types”
- Of interest are
 - » Pupillary Unrest
 - » Fatigue Waves
 - » Pupillary Escape Waves

The Pupil

Afferent Pupillary Defect

1. Confusion with Alpha Omega
2. RAPD due to Unilateral Optic Nerve Neuropathy. Indication of retinal disease or optic nerve pathology
2. Re-dilation referred to in literature as "Pupillary Escape"
3. Normal consensual with subnormal direct. Decreased PLR. Anisocoria at times
4. Swinging Flashlight Test
5. Characteristic observation is "release" or re-dilation of both pupils when the light is moved from the normal to the affected eye.



Afferent Pupillary Defect vs. Normal (Loewenfeld "The Pupil")




Figure 17-4. Pupillary "escape" in a patient with afferent impairment (line 1), compared with a normal subject (lines 2 and 3). The patient was an obese 15-year-old girl who complained of severe headaches. She was found to have bilateral papilledema and increased intracranial pressure. Extensive diagnostic investigation failed to reveal the cause of her trouble. During the next months the working diagnosis was pseudotumor cerebri. She was treated with steroids and the papilledema receded. About 2 months later her headache recurred and her vision was reduced to light perception. One month later a craniotomy was done for decompression, and the cortex and meninges were found to be studded with nodules of metastatic melanoma. The patient died about 2 months later; and since only the brain was examined, the location of the primary tumor remained unknown. The neuropathologist's report stated that "micro-


the patient's reactions to bright light (A) and to darkness (B). Line 2 shows a normal person's reactions under the same conditions. In line 3 light of reduced intensity was used in the normal person (attenuated by neutral grey filter of 10⁻² density). Note the "escape" of the patient's light reflex, compared to the firm, extensive contraction of the normal subject's pupils. With 2 log units of neutral grey filter interposed between the normal eye and the light the same pupillary "escape" was produced in the normal subject as in the patient with bright light. In the darkness reflexes the normal pupil dilated well to one-second interruptions of the light (white bars); and they recontracted quickly when the light was readmitted. In contrast, both the patient's eye and the normal eye with attenuated light had "low intensity" darkness reflexes: the dark-dilations were missing and the secondary contractions were shallow and short-lasting. (From H.S.





The Pupil Reactions

Pupillary Light Reflex



The Pupil Reactions

Near Reflex

1. Observed reaction of Pupil when the patient looks from a distance to near object.
2. Record strength of constriction from 0 – 4+ for each eye.
3. Is a different pathway and can react even if no PLR. Light/Near Dissociation. Afferent pathway problem significantly Dorsal Midbrain.

The Pupil

TESTING

The Pupil TESTING

Conventional

Automated Pupillometry

The Pupil

Testing Standards in Measurement

Observation and Recording

General Observations and Questions (Loewenfeld "The Pupil")

1. Size - too large/small for age, illumination, etc
2. Direct and Consensual responses.
3. Are reactions to light and to near equally extensive
4. Are they equal in size
5. If unequal is difference greater in dim or bright
6. Do both constrict to light
7. Do both redilate as light removed
8. Is reflex to sensory stimuli intact on both sides
9. Are there other motor or sensory defects relating to the pupillary syndrome

The Pupil

Testing Constants in Measurement

Rigid Test Protocol

- Room Illumination
- Patient Fixation
- Light Source
- Distance and Location
- Duration

The Pupil

Testing Constants in Measurement

Room Illumination

- Dimly lit room in dark adapted state

Patient Fixation

- At distance (non-accommodative to avoid near reflex)
- Non-descript target – no cognition

The Pupil

Testing Constants in Measurement

Light Source

- Small, bright, concise, with ability to adjust the light intensity



Pulaski AO Pupil

The Pupil

Testing Constants in Measurement

Distance and Location

- Approximately 6-8" from eye
- Light source turned on and below or temporal to eye being tested
- Light should be swung directly to a location straight into the line of sight
- Lag of 2-3 seconds between eyes



The Pupil

Testing Constants in Measurement

Duration

Normally up to 2-3 seconds in front of eye. Needs to be at least 1 second to get full constriction.

Alpha Omega evaluation

- Observe the pupil under constant light stimulation for at least ten seconds or until first sustained release.
- Test right eye first and then immediately left eye
- Observe time to release, fluctuations and amplitude
- Repeat at least three times observing changes with fatigue

The Pupil TESTING

Automated Pupillometry

Testing - Pupillometry

Conventional

- Subjective with many areas of variability. Slow reactive pupils less than 0.3mm and anisocoria.


Automated Pupillometry

- Reliable, quantitative, objective testing
- Reproducible needed for comparative testing over time especially important in monitoring our treatments and for research.
- Accurate immediate assessment of patient.
- More sensitive with smaller inter-examiner discrepancies.

The Pupil

Testing – Pupillometry

Bright Lamp



The Pupil

Testing – Pupilometry

Bright Lamp


Patient Information

Birth Date	05/09/2008
Sex	Female
Eye Location	Left
Concussion Date	N/A
Asymptomatic?	Yes

Test Information

Test Date & Time	03/12/2020 05:27:57 PM
Test Administrator	John Pulaski/OD
Test Type	Full Spectrum w/Torch
Torch Level	50%
Flash Status	Off
Video Duration	10.5 sec
Torch Duration	10 sec

Pupillogram



Results

Metric	Results (red values are 2σ > mean normative range)	Normative Data
(Beta) Reflex Score	Error: 3	0 - 5
Avg. Constriction Speed	1.762 mm/sec	0.45 - 1.159 mm/sec
Avg. Diameter	4.259 mm	3.169 - 3.816 mm
Avg. Dilation Velocity	N/A	0.888 - 2.699 mm/sec
Constriction Time	2.1 sec	1.178 - 2.217 sec
Latency	0.233 sec	0.138 - 0.279 sec
Max. Constriction Speed	8.4 mm/sec	4.013 - 9.136 mm/sec
Max. Diameter	7.229 mm	3.662 - 5.186 mm
Min. Diameter	3.703 mm	2.861 - 3.317 mm
75% Recover Time	N/A	3.945 - 4.54 sec


Patient Information

Birth Date	03/11/2006
Sex	Male
Eye Location	Left
Concussion Date	N/A
Asymptomatic?	Yes

Test Information

Test Date & Time	03/13/2020 09:10:42 AM
Test Administrator	John Pulaski/OD
Test Type	No Light
Torch Level	0%
Flash Status	Off
Video Duration	10.5 sec
Torch Duration	10 sec

Pupillogram






Results

Metric	Results (red values are 2σ > mean normative range)	Normative Data
Avg. Diameter	3.958 mm	3.626 - 5.161 mm
Max. Diameter	5.02 mm	3.825 - 5.545 mm
Min. Diameter	3.05 mm	3.427 - 4.847 mm

Reflex - PLR Analyzer is designed to help assess and monitor cognitive and neuro-ocular function by providing quantitative data of the Pupillary Light Reflex (PLR). This data can be used by health care professionals to assist in diagnostic decisions and treatment/recovery management. For more details regarding the clinical relevance of PLR, and for more information about Reflex, go to brightlamp.org.

Pupil Diameter






Pulaski AO Pupil

The Pupil

Testing – Bright Lamp



The Pupil

Testing – Bright Lamp

Pupil Parameter Normal Ranges

• Avg. Constriction Speed	0.45 - 1.159 mm/sec
• Avg. Diameter	3.169 - 3.816 mm
• Avg. Dilation Velocity	0.888 - 2.699 mm/sec
• Constriction Time	1.178 - 2.217 sec
• Latency	0.138 - 0.279 sec
• Max. Constriction Speed	4.013 - 9.136 mm/sec
• Max. Diameter	3.662 - 5.186 mm
• Min. Diameter	2.861 - 3.317 mm
• 75% Recovery Time	3.945 - 4.54 sec

The Pupil

Automated Testing – BrightLamp Reflex

The Pupillary Light Reflex as a Biomarker of Concussion

Frederick Robert Carrick,^{1,2,3,4,5,*} Sergio F. Azzolino,⁵ Melissa Hunfalvay,⁵ Guido Pagnacco,^{5,6} Elena Oggero,^{5,6} Ryan C. N. D'Arcy,^{7,8,9} Mahera Abdulrahman,¹⁰ and Kiminobu Sugaya^{1,2}
Gary Peh, Academic Editor

Life (Basel). 2021 Oct; 11(10): 1104.

Published online 2021 Oct 18. doi: [10.3390/life11101104](https://doi.org/10.3390/life11101104)

PMCID: PMC8537991

PMID: [34685475](https://pubmed.ncbi.nlm.nih.gov/34685475/)

The Pupillary Light Reflex as a Biomarker of Concussion

Frederick Robert Carrick et al 2021

- Looked at the parameters of the PLR in concussed versus non-concussed patients, presence of concussion symptoms, age and gender.
- Large Retrospective Study reviewing their clinical patient records – 01/2019 to 01/2020. Over 20,000 patients.
- Used of new automated pupillary testing – BrightLamp Reflex iPhone App. Reproducible diagnostic test.



Questions or Comments?

Matthew Southern
Marketing Director

msouthern@brightlamp.org

Phone: (317) 763-0786



Try Reflex Free For 60-Days:


<https://apps.apple.com/us/app/reflex-plr-analyzer/id1412154869>

The Pupil

REACTIONS OF PUPIL RELATE TO SIZE OF FIELD



The Kinetic Visual Field


The Normal Visual Field



Kinectic (Functional) Visual Fields

Abnormal






5.25 to 3.75 to 3.5 re dilates to 4.25

Alie Koli, xhensiq

4-20-06



The Pupil

Grading Standardization

Alpha Omega Pupil

Alpha Omega Grading Standard
(Pulaski 2010)


Grade	Release Time	Fluctuations	Amplitude
Normal	≥ 7 seconds	Trace	Trace
1+ AO	4 – 6 sec	Moderate	Mild
2+ AO	2 – 3 sec	Marked	Mild-Moderate
3+ AO	1 – 2 sec	Mild-Moderate	Moderate
4+ AO	< 1 sec	Mild	Large



4+ PLR, 5 to 3.0 then release late to 3.75 1-2+AO





OD 6.5 to 4.25 to 4.75






1-2+, 3-4+ PLR 7.0 to 4.5 Release to 5.0







3+AO PLR 2+ 1st 8.0 to 4.16 Release 5.40.
2nd 7.0 to 4.24 to 5.40









AO 4+ 6.8 to 5.5 with 1-2 +PLR



















Pulaski AO Pupil



Thank you!